



**Osborne**  
Co-operative Academy Trust

Self-help  
Self-responsibility  
Equity  
Equality  
Democracy  
Solidarity

## **Infection Prevention and Control Policy**

**First approved by Trust Board: July 2020**

**Review Frequency: Every Two Years**

**Date of last review: April 2022**

**Date of next review: April 2024**

## Version Control

Author	Date Created	Version	Notes
L.Coates	July 2020	2.0	Converted to Trust policy from local school policy
L. Coates	September 2020	3.0	Page 10 – Paragraph added in line with government guidance issued on 17/9/2020 Page 10 – Paragraph added regarding pregnancy during Covid19
L Coates	March 2022	4.0	Page 1- Removal of Covid appendix Page 5 – Links to new Covid guidance added Page 8 – Removal of actions when other members of the household have Covid-19 symptoms Page 11 – Covid-19 guidance link added Page13 – Covid -19 update to actions taken in the event of positive case
L Coates	April 2022	5.0	Page 9 – Testing positive for Covid-19 Page 10 – Statement of actions taken for Covid-19 symptoms Page 11 – PHE guidance <a href="#">Respiratory Infections, including Covid-19</a> Page 13 – PHE guidance on absence following positive test

## **Contents:**

### [Statement of intent](#)

Preventative measures

[Legal framework](#)

[Ensuring a clean environment](#)

[Pupil immunisation](#)

[Staff immunisation](#)

[Contact with pets and animals](#)

[Water-based activities](#)

In the event of infection

[Preventing the spread of infection](#)

[Vulnerable pupils](#)

[Procedures for unwell pupils/staff](#)

[Exclusion](#)

Medication

[Outbreaks of infectious diseases](#)

Pregnant staff members

[Staff handling food](#)

[Managing specific infectious diseases](#)

[Monitoring and review](#)

Appendices

Managing Specific Infectious Diseases

[Infection Absence Periods](#)

[Diarrhoea and Vomiting Outbreak Action Checklist](#)

[List of Notifiable Diseases](#)

## Statement of intent

Infections can easily spread in a school due to:

- Pupils' immature immune systems.
- The close-contact nature of the environment.
- Some pupils having not yet received full vaccinations.
- Pupils' poor understanding of good hygiene practices.

Infections commonly spread in the following ways:

- Respiratory spread – contact with coughs or other secretions from an infected person.
- Direct contact spread – direct contact with the infecting organism, e.g., skin-on-skin contact during sports.
- Gastrointestinal spread – contact with contaminated food or water, or contact with infected faeces or unwashed hands.
- Blood borne virus spread – contact with infected blood or bodily fluids, e.g., via bites or used needles.

We actively prevent the spread of infection via the following measures:

- Maintaining high standards of personal hygiene and practice
- Maintaining a clean environment
- Routine immunisations
- Taking appropriate action when infection occurs, including additional cleaning routines

This policy aims to help school staff prevent and manage infections in school. It is not intended to be used as a tool for diagnosing disease, but rather a series of procedures informing staff what steps to take to prevent infection and what actions to take when infection occurs.

Signed by



Headteacher

Date 29<sup>th</sup> April 2022



Chair of Governors

Date 29<sup>th</sup> April 2022

## Legal framework

This policy has due regard to legislation including, but not limited to, the following:

- Control of Substances Hazardous to Health Regulations 2002 (as amended 2004)
- Health and Safety at Work etc. Act 1974
- The Management of Health and Safety at Work Regulations 1999
- The Reporting of Injuries, Diseases and Dangerous Occurrences Regulations 2013
- The Health Protection (Notification) Regulations 2010

This policy has due regard to statutory guidance including, but not limited to, the following:

- Public Health England (2019) 'Health protection in schools and other childcare facilities'
- DfE (2015) 'Supporting pupils at school with medical conditions'
- [Covid-19 Response- Living with Covid](#)

This policy operates in conjunction with the following school policies and documents:

- Incident Management Plan
- Pandemic Incidents Policy
- Health and Safety Policy
- Supporting Pupils with Medical Conditions Policy
- Administering Medication Policy
- First Aid Policy
- Visit to the Zoo Risk Assessment
- Farm Visit Risk Assessment
- Swimming Risk Assessment
- Sharps Policy
- Disposal of Bodily Fluid Policy to be implemented

## Preventative measures

Ensuring a clean environment

Sanitary facilities

- Wall-mounted soap dispensers are used in all toilets – bar soap is never used.
- Waste paper bin is always made available where disposable paper towels are used.
- Toilet paper is always available in cubicles.
- Suitable sanitary disposal facilities are provided where necessary.

Personal Care changing areas

Pupils are encouraged to clean with disposable wipes where their medical needs require, and lotions are labelled with the relevant pupil's name

- Handwashing facilities are available in the room and soiled nappies are disposed of inside a wrapped plastic bag and placed in a nappy dispenser.

## Continence aid facilities

- Pupils who use continence aids, e.g., continence pads and catheters are encouraged to be as independent as possible. Pads are changed in a designated area with adequate handwashing facilities, and disposable powder-free latex gloves and a disposable plastic apron are worn by the adult supporting.

## Laundry

- Manual sluicing of clothing is not permitted and gloves and aprons are worn when handling soiled linen or clothing. Hands are thoroughly washed after gloves are removed.

## Cleaning contractors/School Cleaners

- Both school employed cleaners and cleaning contractor are employed to carry out rigorous cleaning of the premises. Cleaning equipment is maintained to a high standard and is colour coded according to area of use. The school Site Manager/Caretaker is responsible for monitoring cleaning standards and discussing any issues that may arise with the contractor or addresses issues arising with the school cleaners where appropriate.

## Toys and equipment

### Handwashing

- All staff and pupils are advised to wash their hands after using the toilet, before eating or handling food, following outdoor activities and after touching animals.
- Sanitiser dispensers are located at various points around the school.

### Blood and other bodily fluids

- Cuts and abrasions are covered with waterproof dressings.
- When coughing or sneezing, all staff and pupils are encouraged to cover their nose and mouth with a disposable tissue and dispose of the tissue after use, and to wash their hands afterwards.
- Personal protective equipment (PPE) is worn where there is a risk of contamination with blood or bodily fluids during an activity. Gloves are disposable, non-powdered vinyl or latex (staff to be responsible in advising if they are intolerant to latex) and CE (Conformité Européene) marked. If there is a risk of splashing to the face, goggles are worn.
- All staff receive training in Donning and Doffing of PPE.
- Spillages of blood, faeces, saliva, vomit, nasal and eye discharges are cleaned up immediately by the attending adult in personal or medical care and site staff in response to vomiting. They are cleaned using a mixture of detergent and disinfectant. Paper towels or cloths are used, staff always wearing PPE, and they are disposed of after use. **The school spillage kit is stored in the front office.**

### Bites

- If a bite does not break the skin, the affected area is cleaned with soap and water.
- If a bite breaks the skin, the affected area is cleaned with soap and running water, the incident is recorded in the pupil accident log and medical advice is sought immediately.

### Hypodermic needles (sharps)

- Injuries incurred through sharps found on school grounds will be treated in line with Havering oneSource Guidelines. All sharps found on school premises will be disposed of in the sharps bin wearing PPE.

## Pupil immunisation

- The school keeps up-to-date with national and local immunisation scheduling and advice via [www.nhs.uk/conditions/vaccinations/](http://www.nhs.uk/conditions/vaccinations/).

Whilst the school encourages parents/carers to have their children immunised, parental consent will always be sought before a vaccination is given.

- The school will ensure that any pupils with existing medical conditions are medically cleared to be given the vaccine in question.
- A healthcare team will visit the school in order to carry out vaccinations
- A risk assessment will be conducted by the healthcare team prior to vaccinations
- Before starting school, pupils should be given their 4-in-1 pre-school booster against diphtheria, tetanus, whooping cough and polio, usually at 3 years and 4 months.
- All pupils in Reception to Year 4 will be offered nasal flu vaccinations annually.
- Any pupils who become unwell after receiving a vaccination will be treated by following the school's procedures for sick and unwell pupils.
- [COVID-19 vaccination programme for children and young people: guidance for schools \(version 3\) - GOV.UK \(www.gov.uk\)](https://www.gov.uk/guidance/covid-19-vaccination-programme-for-children-and-young-people-guidance-for-schools-version-3)

Any side effects from the vaccinations, such as becoming unwell, will be reported to the healthcare team who administered the vaccination, allowing them to record the symptoms and the time that the vaccine was administered.

- Regular communication is maintained after pupils return to lessons, as some side effects can take several hours to develop.
- Members of staff will be with pupils before, during and after vaccinations, in order to keep the pupils relaxed and create a calming atmosphere.
- The school will ensure that the venue used is a clean, open, well-ventilated room, where pupils can access water and fresh air.
- Needles are kept away from pupils before and after the vaccine is administered.
- Some vaccinations may involve an exclusion period in which pupils are not required to attend school. The administering healthcare team will provide advice in such cases.

### Staff immunisation

Staff are encouraged to remain up-to-date with immunisations; in particular, we encourage the following:

- Hepatitis B: We do not recommend Hepatitis B vaccines for staff in routine contact with infected children; however, where staff are involved with the care of children with severe learning disabilities or challenging behaviour, we encourage immunisation.
- Rubella: Female staff of childbearing age are encouraged to check with their GP that they are immune to the rubella (German measles) virus. If they are not immune, we encourage them to be immunised with the MMR vaccine, except during pregnancy.
- [COVID-19 vaccination programme for children and young people: guidance for schools \(version 3\) - GOV.UK \(www.gov.uk\)](https://www.gov.uk/guidance/covid-19-vaccination-programme-for-children-and-young-people-guidance-for-schools-version-3)

### Contact with pets and animals

- Visiting animals for the purpose of education in school are only permitted in the following areas: **the playground and halls** following a risk assessment.
- Visiting animals are always supervised by their handler, following any physical contact, hands will be washed.
- Visits to farms are strictly controlled by the policies and protocols contained in our Farm Visit Risk Assessment.
- Visits to zoos are strictly controlled by use of our Visit to the Zoo Risk Assessment.

## Water-based activities

### Swimming lessons

- General swimming lessons are governed by the control measures outlined in our Swimming Risk Assessment.
- Pupils who have experienced vomiting or diarrhoea in the week preceding the trip are not permitted to attend public swimming pools.

### Other activities

- Alternative water-based activities are only undertaken at reputable centres.
- Children and staff cover all cuts, scratches and abrasions with waterproof dressings before taking part, and hands are washed immediately after the activity. No food or drink is to be consumed until hands have been washed.
- After canoeing or rowing, staff and pupils immediately wash or shower.
- If a member of staff or a pupil becomes ill within three to four weeks of an activity taking place, we encourage them to seek medical advice and inform their GP of their participation in these activities.

## In the event of infection

### Preventing the spread of infection

Parents/Carers will not bring their child to school in the following circumstances:

- The child shows signs of being poorly and needing one-to-one care
- The child has untreated conjunctivitis
- The child has a high temperature/fever
- The child has untreated head lice
- The child has been vomiting and/or had diarrhoea within the last 48 hours due to illness
- The child has an infection and the [minimum recommended exclusion period](#) has not yet passed
- During or following a pandemic government guidance should be followed for attending school

### Vulnerable pupils

- Pupils with impaired immune defence mechanisms (known as immune-compromised) are more likely to acquire infections. In addition, the effect of an infection is likely to be more significant for such pupils. These pupils may have a disease that compromises their immune system or be undergoing treatment, such as chemotherapy, that has a similar effect.
- Parents/Carers are responsible for notifying the school if their child is “vulnerable”.
- If a vulnerable child is thought to have been exposed to an infectious disease, the child’s parents/carers will be informed and encouraged to seek medical advice from their doctor or specialist.

### Vulnerable Staff

- Appropriate action for staff considered at high risk due to personal medical grounds or caring responsibilities will be taken into account on a case by case basis on a local level. This could include changes to working practices to minimise the risk of infection.

## Procedures for unwell pupils/staff

Staff are required to know the warning signs of pupils becoming unwell including, but not limited to, the following:

- Not being themselves
- Not having a snack
- Not eating at lunchtimes
- Wanting more attention/sleep than usual
- Displaying physical signs of being unwell, e.g., watery eyes, a flushed face or clammy skin
- Lack of senses for taste or smell
- Any symptoms linked to a current pandemic

Where a staff member identifies a pupil as unwell, the pupil is taken to the medical room near the main reception office, where their temperature will be taken by the first aider, and the pupil's parents/carers will be informed of the situation.

The First Aider will:

- Follow hand hygiene guidance before donning and following doffing of PPE
- Don appropriate PPE following the PHE guidance for donning and doffing
- Attempt to cool the pupil down if they are too hot, by opening a window and suggesting that the pupil removes their top layers of clothing.
- Provide the pupil with a drink of water.
- Move the pupil to a quieter area of the classroom or school such as the medical room.
- Ensure there is a staff member available to comfort the pupil.
- Summon emergency medical help if required.

Pupils and staff displaying any of the signs of becoming unwell will be sent home, and we will recommend that they see a doctor.

If a pupil is identified with sickness and diarrhoea, the pupil's parents/carers will be contacted immediately and the child will be sent home and may only return after 48 hours have passed without symptoms if this was due to illness.

If a staff member is suffering from vomiting and diarrhoea, they will be sent home and may not return until 48 hours have passed without symptoms.

In the event of testing positive for Covid-19, pupils and staff are advised to stay away from school for five days, staff should then have a negative lateral flow test (LFT) prior to returning to work. If an LFT is not available staff should only return if they no longer have a temperature and they feel well enough.

If the school is unable to contact a pupil's parents/carers in any situation, the pupil's alternative emergency contacts will be contacted.

### **Contaminated clothing**

If the clothing of the first-aider or a pupil becomes contaminated, the clothing is removed as soon as possible and placed in a plastic bag. The pupil's clothing is sent home with the pupil.

### **Exclusion**

- Pupils suffering from infectious diseases will be excluded from school on medical grounds for the minimum recommended period.
- Pupils can be formally excluded on medical grounds by the Head of School.

- If parents/carers insist on their child returning to school when the child still poses a risk to others, following advice from PHE or the HPT, the Trust may serve notice on the child's parents/carers to require them to keep the child away from school until the child no longer poses a risk of infection.
- If a pupil is exposed to an infectious disease, but is not confirmed to be infected, this is not normally a valid reason for exclusion; however, the local health protection team (HPT) through the Eastern England Health Protection Team via 03003038537 may be contacted to advice on a case-by-case basis.

## Medication

- Where a pupil has been prescribed medication by a doctor, dentist, nurse or pharmacist, the first dose will be given at home, in case the pupil has an adverse reaction.
- The pupil will only be allowed to return to school 24 hours after the first dose of medication, to allow it time to take effect.
- All medicine provided in school will be administered in line with the Administering Medication Policy.

## Outbreaks of infectious diseases

**In the event of a pandemic we will follow the guidance set out by Public Health England and the DfE and invoke our Incident Management Plan and the Pandemic Incidents Policy.**

An incident is classed as an 'outbreak' where:

- Two or more people experiencing a similar illness are linked in time or place.
- A greater than expected rate of infection is present compared with the usual background rate, e.g.:
- Two or more pupils in the same classroom are suffering from vomiting and diarrhoea.
- A greater number of pupils than usual are diagnosed with scarlet fever.
- There are two or more cases of measles at the school.
- Schools will manage local outbreaks of COVID-19 as they do with other respiratory infection, inform parents to be vigilant and ensure cleaning regime is adhered to and there is good ventilation maintained in the learning areas.

Schools will Inform the Trust if there is:

- evidence of severe disease due to COVID-19, for example if a pupil, student, or staff member is admitted to hospital due to COVID-19
- a cluster of cases where there are concerns about the health needs of vulnerable staff or students within the affected group

Settings can seek public health and operational advice by phoning the DfE helpline (0800 046 8687, option 1), or ring Thurrock PH Team on [01375 652510](tel:01375652510)

Respiratory hygiene The 'catch it, bin it, kill it' approach continues to be very important. [The e-Bug COVID-19 website](#) contains free resources for you, including materials to encourage good hand and respiratory hygiene

Suspected outbreaks of any of the diseases listed on the [List of Notifiable Diseases](#) will always be reported.

As soon as an outbreak is suspected (even if it cannot be confirmed), the Head of School will contact the HPT by calling: 03003038537 Eastern England Health Protection Team to discuss the situation and agree if any actions are needed.

During an outbreak, enhanced cleaning protocols will be undertaken, following advice provided by the local HPT. The Site Manager/Caretaker will liaise with the cleaning contractor to ensure these take place.

## **Pregnant staff members**

If a pregnant staff member develops a rash or is in direct contact with someone who has a potentially contagious rash, we will strongly encourage her to speak to her doctor or midwife.

## **Covid19 Pandemic**

Pregnancy risk assessments will be reviewed each trimester recognising the Coronavirus as a hazard and implementing control measures to ensure the safety of staff. As a vulnerable group, individual risk assessments will be completed with agreement from the member of staff using the oneSource tool and a copy placed on the personal file. The school will follow guidance from the government, NHS, HSE and HR advisors to protect the well-being of pregnant staff.

## **Chickenpox**

If a pregnant staff member has not already had chickenpox or shingles, becoming infected can affect the pregnancy. If a pregnant staff member believes they have been exposed to chickenpox or shingles and have not had either infection previously, she will speak to her midwife or GP as soon as possible. If a pregnant staff member is unsure whether they are immune, we encourage them to take a blood test.

## **Measles**

If a pregnant staff member is exposed to measles, she will inform her midwife immediately.

## **Rubella (German measles)**

If a pregnant staff member is exposed to rubella, she will inform her midwife immediately.

## **Respiratory Infection including coronavirus (COVID-19)**

### [Guidance from Public Health England](#)

Respiratory infections are common in children and young people, particularly during the winter months. Symptoms can be caused by several respiratory infections including the common cold, COVID-19, flu, and respiratory syncytial virus (RSV). For most children these illnesses will not be serious, and they soon recover.

Children with respiratory infections can experience a range of symptoms including a runny nose, high temperature, cough and sore throat. It is not possible to tell which germ someone is infected with based on symptoms alone.

Respiratory infections can spread easily between people. Sneezing, coughing, singing and talking may spread respiratory droplets from an infected person to someone close by. Droplets from the mouth or nose may also contaminate hands, eating and drinking utensils, toys or other items and spread to those who may use or touch them, particularly if they then touch their nose or mouth.

Children with mild symptoms such as a runny nose, sore throat, or mild cough, who are otherwise well, can continue to attend their education or childcare setting.

Children and young people who are unwell and have a [high temperature](#) should stay at home and where possible avoid contact with other people. They can go back to education or childcare setting when they no longer have a high temperature and they are well enough.

### **Slapped cheek disease (Parvovirus B19)**

If a pregnant staff member is exposed to slapped cheek disease, she will inform her midwife promptly.

### **Staff handling food**

Food handling staff suffering from transmittable diseases will be excluded from all food handling activity until advised by the Catering Company serving the school that they are clear to return to work. Both food handling staff and midday assistants are not permitted to attend work if they are suffering from diarrhoea and/or vomiting. They are not permitted to return to work until 48 hours have passed since diarrhoea and/or vomiting occurred.

Food handlers are required by law to inform the school if they are suffering from any of the following:

- Typhoid fever
- Paratyphoid fever
- Other salmonella infections
- Dysentery
- Shigellosis
- Diarrhoea (where the cause of which has not been established)
- Infective jaundice
- Staphylococcal infections likely to cause food poisoning like impetigo, septic skin lesions, exposed infected wounds, boils
- E. coli VTEC infection

### **Managing specific infectious diseases**

When an infectious disease occurs in the school, we will follow the appropriate procedures set out in the Managing Specific Infectious Diseases appendix.

### **Monitoring and review**

All members of staff are required to familiarise themselves with this policy as part of their induction programme.

The Head of School will review this policy on an annual basis and will make any changes necessary, taking into account the current effectiveness of infection control and prevention.

The next scheduled review date is March 2023 or in the event of changes to the Pandemic guidance.

## Managing Specific Infectious Diseases

Disease	Symptoms	Considerations	Exclusion period
Athlete's foot	Scaling or cracking of the skin, particularly between the toes, or blisters containing fluid. The infection may be itchy.	Cases are advised to see their GP for advice and treatment.	Exclusion is not necessary.
Chicken pox	Sudden onset of fever with a runny nose, cough and generalised rash. The rash then blisters and scabs over. Several blisters may develop at once, so there may be scabs in various stages of development. Some mild infections may not present symptoms.	Cases are advised to consult their GP.	Chickenpox is infectious from 48 hours prior to a rash appearing up to five days after the onset of a rash.  Cases will be excluded from school for five days from the onset of a rash.  It is not necessary for all the spots to have healed before the case returns to school.
Cold sores	The first signs of cold sores are tingling, burning or itching in the affected area. Around 24 hours after the first signs appear the area will redden and swell, resulting in a fluid-filled blister. After blistering, they break down to form ulcers then dry up and crust over.	Cases are advised not to touch the cold sore, or to break or pick the blisters. Sufferers of cold sores should avoid kissing people and should not share items such as cups, towels and facecloths.	Exclusion is not necessary.
Conjunctivitis	The eye(s) become reddened and swollen, and there may be a yellow or green discharge. Eyes may feel itchy and 'gritty'.	Cases are encouraged to seek advice, wash their hands frequently and not to rub their eyes.  The HPT will be contacted if an outbreak occurs.	Exclusion is not necessary.
Coronavirus	A new, continuous cough and a high temperature or a loss or change to your sense of smell or taste are the main symptoms of coronavirus. Most people with Coronavirus have at least one of these symptoms.	In the Osborne Co-operative Academy Trust any staff or pupils with symptoms of a respiratory infection, such as COVID-19 which includes having a high temperature, or you do not feel well enough to go to work/school or carry out normal activities, you are advised to try to stay at home and avoid contact with other people.	The usual sickness absence guidance will be followed unless guidance from the government states otherwise.  If staff or pupils choose to take a test and it is positive, <a href="#">PHE guidance</a> on exclusion should be followed.

Disease	Symptoms	Considerations	Exclusion period
Food poisoning	Symptoms normally appear within one to two days of contaminated food being consumed, although they may start at any point between a few hours and several weeks later. The main symptoms are likely to be nausea, vomiting, diarrhoea, stomach cramps and fever.	<p>Cases will be sent home.</p> <p>The HPT will be contacted where two or more cases with similar symptoms are reported.</p> <p>The cause of a food poisoning outbreak will always be investigated.</p>	<p>Cases will be excluded until 48 hours have passed since symptoms were present.</p> <p>For some infections, longer exclusion periods may be required. The HPT will advise in such cases.</p>
Giardia	Symptoms include abdominal pain, bloating, fatigue and pale, loose stools.	<p>Cases will be sent home.</p> <p>The HPT will be contacted where two or more cases with similar symptoms are reported.</p>	Cases will be excluded until 48 hours have passed since symptoms were present.
Salmonella	Symptoms include diarrhoea, headache, fever and, in some cases, vomiting.	<p>Cases will be sent home.</p> <p>The HPT will be contacted where two or more cases with similar symptoms are reported.</p>	Cases will be excluded until 48 hours have passed since symptoms were present.
Typhoid and paratyphoid fever	Symptoms include tiredness, fever and constipation. The symptoms of paratyphoid fever include fever, diarrhoea and vomiting.	All cases will be immediately reported to the HPT.	<p>Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved.</p> <p>Environmental health officers or the HPT may advise the school to issue a lengthened exclusion period.</p>
E. coli (verocytotoxigenic or VTEC)	Symptoms vary but include diarrhoea, abdominal cramps, headaches and bloody diarrhoea.	Cases will immediately be sent home and advised to speak to their GP.	<p>Cases will be excluded whilst symptomatic and for 48 hours after symptoms have resolved.</p> <p>Where the sufferer poses an increased risk, for example, food handlers, they will be</p>

Disease	Symptoms	Considerations	Exclusion period
			<p>excluded until a negative stool sample has been confirmed.</p> <p>The HPT will be consulted in all cases.</p>
Gastroenteritis	Symptoms include three or more liquid or semi-liquid stools in a 24-hour period.	The HPT will be contacted where there are more cases than usual.	<p>Cases will be excluded until 48 hours have passed since symptoms were present.</p> <p>If medication is prescribed, the full course must be completed and there must be no further symptoms displayed for 48 hours following completion of the course before the cases may return to school.</p> <p>Cases will be excluded from swimming for two weeks following their last episode of diarrhoea.</p>
Bacillary dysentery (Shigella)	Symptoms include bloody diarrhoea, vomiting, abdominal pain and fever. It lasts four to seven days on average, but potentially several weeks.	The school will contact the HPT.	Microbiological clearance is required for some types of shigella. The HPT will advise.
Campylobacter	Symptoms include diarrhoea, headache, fever and, in some cases, vomiting.		Cases will be excluded until 48 hours have passed since symptoms were present.
Cryptosporidiosis	Symptoms include abdominal pain, diarrhoea and occasional vomiting.		Cases will be excluded until 48 hours have passed since symptoms were present.
Glandular fever	Symptoms include severe tiredness, aching muscles, sore throat, fever, swollen glands and occasionally jaundice.	The sufferer may feel unwell for several months and the school will provide reasonable adjustments where necessary.	Exclusion is not necessary, and cases can return to school as soon as they feel well.
Hand, foot and mouth disease	Symptoms include a fever, reduced appetite and generally feeling unwell. One or two days later, a rash with blisters will develop on cheeks, hands and feet. Not all cases will have symptoms.		Exclusion is not necessary, and cases can return to school as soon as they feel well.
Head lice	Other than the detection of live lice or nits, there are no immediate symptoms until two to three weeks	Treatment is only necessary when live lice are seen.	Exclusion is not necessary.

Disease	Symptoms	Considerations	Exclusion period
	after infection, where itching and scratching of the scalp occurs.	<p>Staff are not permitted to inspect any pupil's hair for head lice.</p> <p>If a staff member incidentally notices head lice in a pupil's hair, they will inform the pupil's parents and advise them to treat their child's hair.</p> <p>When a pupil has been identified as having a case of head lice, a letter will be sent home to all parents notifying them that a case of head lice has been reported and asking all parents to check their children's hair.</p>	
Hepatitis A	Symptoms include abdominal pain, loss of appetite, nausea, fever and tiredness, followed by jaundice, dark urine and pale faeces.	The illness in children usually lasts one to two weeks but can last longer and be more severe in adults.	<p>Cases are excluded while unwell and for seven days after the onset of jaundice (or the onset of symptoms if no jaundice presents), the case is under five years of age or where hygiene is poor.</p> <p>There is no need to exclude older children with good hygiene.</p>
Hepatitis B	Symptoms include general tiredness, nausea, vomiting, loss of appetite, fever and dark urine, and older cases may develop jaundice.	<p>The HPT will be contacted where advice is required.</p> <p>The procedures for dealing with blood and other bodily fluids will always be followed.</p> <p>The accident book will always be completed with details of injuries or adverse events related to cases.</p>	<p>Acute cases will be too ill to attend school and their doctor will advise when they are fit to return.</p> <p>Chronic cases will not be excluded or have their activities restricted.</p> <p>Staff with chronic hepatitis B infections will not be excluded.</p>

Disease	Symptoms	Considerations	Exclusion period
Hepatitis C	Symptoms are often vague but may include loss of appetite, fatigue, nausea and abdominal pain. Less commonly, jaundice may occur.	The procedures for dealing with blood and other bodily fluids will always be followed.  The accident book will always be completed with details of injuries or adverse events related to cases.	Cases will not be excluded or have their activities restricted.
Impetigo	Symptoms include lesions on the face, flexures and limbs.	Towels, facecloths and eating utensils will not be shared by pupils.  Toys and play equipment will be cleaned thoroughly.	Cases will be excluded until lesions have healed and crusted or 48 hours after commencing antibiotic treatment.
Influenza	Symptoms include headache, fever, cough, sore throat, aching muscles and joints, and tiredness.	Those in risk groups will be encouraged to have the influenza vaccine.  Anyone with flu-like symptoms will stay home until they have recovered. Pupils under 16 will not be given aspirin.	Cases will remain home until they have fully recovered.
Measles	Symptoms include a runny nose, cough, conjunctivitis, high fever and small white spots around the cheeks. Around the third day, a rash of flat red or brown blotches may appear on the face then spread around the body.	All pupils are encouraged to have MMR immunisations in line with the national schedule.  Staff members should be up to date with their MMR vaccinations.  Pregnant staff members and those with weak immune systems will be encouraged to contact their GP immediately for advice if they meet measles.	Cases are excluded for four days after the onset of a rash.
Meningitis	Symptoms include fever, severe headaches, photophobia, stiff neck, non-blanching rash, vomiting and drowsiness.	Meningitis is a notifiable disease.	Once a case has received any necessary treatment, they can return to school.

Disease	Symptoms	Considerations	Exclusion period
Meningococcal meningitis and meningitis septicaemia	Symptoms include fever, severe headaches, photophobia, stiff neck and a non-blanching rash.	<p>Medical advice will be sought immediately.</p> <p>The confidentiality of the case will always be respected.</p> <p>The HPT and school health advisor will be notified of a case of meningococcal disease in the school. The HPT will conduct a risk assessment and organise antibiotics for household and close contacts.</p> <p>The HPT will be notified if two cases of meningococcal disease occur in the school within four weeks.</p>	<p>When the case has been treated and recovered, they can return to school.</p> <p>Exclusion is not necessary for household or close contacts unless they have symptoms suggestive of meningococcal infection.</p>
Meningitis (viral)	Symptoms include headache, fever, gastrointestinal or upper respiratory tract involvement and, in some cases, a rash.	<p>The case will be encouraged to consult their GP.</p> <p>If more than once case occurs, the HPT will be consulted.</p>	No exclusion is required.
Meticillin resistant staphylococcus aureus (MRSA)	Symptoms are rare but include skin infections and boils.	All infected wounds will be covered.	No exclusion is required.
Mumps	Symptoms include a raised temperature and general malaise. Then, stiffness or pain in the jaws and neck is common. Following this, the glands in the cheeks and under the jaw swell up and cause pain (this can be on one or both sides). Mumps may also cause swelling of the testicles.	<p>The case will be encouraged to consult their GP.</p> <p>Parents are encouraged to immunise their children against mumps.</p>	Cases can return to school five days after the onset of swelling if they feel able to do so.
Ringworm	Symptoms vary depending on the area of the body affected.	Pupils with ringworm of the feet will wear socks and trainers at all times and cover their feet during physical education.	No exclusion is usually necessary.

Disease	Symptoms	Considerations	Exclusion period
			For infections of the skin and scalp, cases can return to school once they have received treatment.
Rotavirus	Symptoms include severe diarrhoea, stomach cramps, vomiting, dehydration and mild fever.	Cases will be sent home if unwell and encouraged to speak to their GP.	Cases will be excluded until 48 hours have passed since symptoms were present.
Rubella (German Measles)	Symptoms are usually mild, with a rash being the first indication. There may also be mild catarrh, headaches or vomiting. There may be a slight fever and some tenderness in the neck, armpits or groin, and there may be joint pains.	MMR vaccines are promoted to all pupils.	Cases will be excluded for six days from the appearance of the rash.
Scabies	Symptoms include tiny pimples and nodules on a rash, with burrows commonly seen on the wrists, palms, elbows, genitalia and buttocks.	All household contacts and any other very close contacts should have one treatment at the same time as the second treatment of the case.  The second treatment must not be missed and should be carried out one week after the first treatment.	Cases will be excluded until after the first treatment has been carried out.
Scarlet Fever	Symptoms include acute inflammation of the pharynx or tonsils, with tonsils reddening in colour and becoming partially covered with a thick, yellowish exudate. In severe cases, there may be a high fever, difficulty swallowing and tender, enlarged lymph nodes. A rash develops on the first day of fever and is red, generalised, pinhead in size and gives the skin a sandpaper-like texture, with the tongue developing a strawberry-like appearance.	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered.  If two or more cases occur, the HPT will be contacted.	Cases are excluded for 24 hours following appropriate antibiotic treatment.
Slapped cheek syndrome, Parvovirus B19, Fifth's Disease	Where symptoms develop, they include a rose-red rash making the cheeks appear bright red.	Cases will be encouraged to visit their GP.	Exclusion is not required.

Disease	Symptoms	Considerations	Exclusion period
Threadworm	Symptoms include itching around the anus, particularly at night.	Cases will be encouraged to visit their GP.	Exclusion is not required.
Tuberculosis (TB)	Symptoms include cough, loss of appetite, weight loss, fever, sweating (particularly at night), breathlessness and pains in the chest. TB in parts of the body other than the lungs may produce a painful lump or swelling.	Advice will be sought from the HPT before taking any action, and regarding exclusion periods.	<p>Cases with infectious TB can return to school after two weeks of treatment if well enough to do so, and as long as they have responded to anti-TB therapy.</p> <p>Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses, will not be excluded.</p>
Whooping cough (pertussis)	Symptoms include a heavy cold with a persistent cough. The cough generally worsens and develops the characteristic 'whoop'. Coughing spasms may be worse at night and may be associated with vomiting.	Cases will be advised to see their GP. Parents are advised to have their children immunised against whooping cough.	<p>Cases will not return to school until they have had 48 hours of appropriate treatment with antibiotics and feel well enough to do so, or 21 days from the onset of illness if no antibiotic treatment is given.</p> <p>Cases will be allowed to return in the above circumstances, even if they are still coughing.</p>

### Infection Absence Periods

This table details the minimum required period for staff and pupils to stay away from school following an infection, as recommended by Public Health England.

\*Identifies a notifiable disease. It is a statutory requirement that doctors report these diseases to their local Public Health England centre.

Infection	Recommended minimum period to stay away from school	Comments
Athlete's foot	None	Treatment is recommended; however, this is not a serious condition.
Chicken pox	Until all vesicles have crusted over	Follow procedures for vulnerable children and pregnant staff.
Cold sores	None	Avoid contact with the sores.
Conjunctivitis	None	If an outbreak occurs, consult the HPT.
Coronavirus	Five days. Staff should have a negative lateral flow test prior to their return, in the absence of this they should only return once they no longer have a temperature and feel well enough.	If an outbreak occurs, consult the HPT.
Diarrhoea and/or vomiting	Whilst symptomatic and 48 hours from the last episode	GPs should be contacted if diarrhoea or vomiting occur after taking part in water-based activities.
Diphtheria*	Exclusion is essential.	Family contacts must be excluded until cleared by the HPT and the HPT must always be consulted.

Infection	Recommended minimum period to stay away from school	Comments
Flu (influenza)	Until recovered	Report outbreaks to the HPT.
Glandular fever	None	
Hand foot and mouth	None	Contact the HPT if a large number of children are affected. Exclusion may be considered in some circumstances.
Head lice	None	Treatment recommended only when live lice seen.
Hepatitis A*	Seven days after onset of jaundice or other symptoms	If it is an outbreak, the HPT will advise on control measures.
Hepatitis B*, C* and HIV	None	Not infectious through casual contact. Procedures for bodily fluid spills must be followed.
Impetigo	48 hours after commencing antibiotic treatment, or when lesions are crusted and healed	Antibiotic treatment is recommended to speed healing and reduce the infectious period.
Measles*	Four days from onset of rash	Preventable by vaccination (MMR). Follow procedures for vulnerable children and pregnant staff.
Meningococcal meningitis*/ septicaemia*	Until recovered	Meningitis ACWY and B are preventable by vaccination.  The HPT will advise on any action needed.

Infection	Recommended minimum period to stay away from school	Comments
Meningitis* due to other bacteria	Until recovered	Hib and pneumococcal meningitis are preventable by vaccination. The HPT will advise on any action needed.
Meningitis viral*	None	As this is a milder form of meningitis, there is no reason to exclude those who have been in close contact with infected persons.
MRSA	None	Good hygiene, in particular environmental cleaning and handwashing, is important to minimise the spread. The local HPT should be consulted.
Mumps*	Five days after onset of swelling	Preventable by vaccination with two doses of MMR.
Ringworm	Exclusion is not usually required	Treatment is required.
Rubella (German measles)	Four days from onset of rash	Preventable by two doses of immunisation (MMR). Follow procedures for pregnant staff.
Scarlet fever	24 hours after commencing antibiotic treatment	Antibiotic treatment is recommended, as a person is infectious for two to three weeks if antibiotics are not administered. If two or more cases occur, the HPT should be contacted.
Scabies	Can return to school after first treatment	The infected person's household and those who have been in close contact will also require treatment.
Slapped cheek/Fifth disease/Parvo Virus B19	None (once rash has developed)	Follow procedures for vulnerable children and pregnant staff.
Threadworms	None	Treatment recommended for the infected person and household contacts.
Tonsillitis	None	There are many causes, but most causes are virus-based and do not require antibiotics.
Tuberculosis (TB)	Pupils with infectious TB can return to school after two weeks of treatment if well enough to do so, and as long as	Only pulmonary (lung) TB is infectious. It requires prolonged close contact to spread. Cases with non-pulmonary TB, and cases with pulmonary TB who have effectively completed two weeks of treatment as confirmed by TB nurses, should not be

Infection	Recommended minimum period to stay away from school	Comments
	they have responded to anti-TB therapy.	excluded. Consult the local HPT before disseminating information to staff and parents.
Warts and verrucae	None	Verrucae should be covered in swimming pools, gymnasiums and changing rooms.
Whooping cough (pertussis)*	Two days from commencing antibiotic treatment, or 21 days from the onset of illness if no antibiotic treatment is given	Preventable by vaccination. Non-infectious coughing can continue for many weeks after treatment. The HPT will organise any necessary contact tracing.

## Diarrhoea and Vomiting Outbreak Action Checklist

Date:	
Completed by:	

Action	Action taken?		Comments
	Yes	No	
A 48-hour exclusion rule has been enforced.			
Liquid soap and paper hand towels or hand dryer are available.			
Enhanced cleaning is undertaken twice daily, and an appropriate disinfectant is used.			
Appropriate personal protective equipment (PPE) is available.			
Appropriate waste disposal systems are available for removing infectious waste.			
Toys are cleaned and disinfected on a regular basis.			
Visitors are restricted, and essential visitors are informed of the outbreak.			
New children joining the school are delayed from joining.			
The health protection team (HPT) has been informed of any infected food handlers.			
Staff work in dedicated areas and food handling is restricted.			
All staff (including agency) are asked if they are unwell.			
Staff are restricted from working elsewhere.			
The HPT is informed of any planned events at the school.			
Ofsted are informed if necessary (Planned Inspection due)			

## List of Notifiable Diseases

Under the Health Protection (Notification) Regulations 2010, the following diseases will always be reported to the health protection team (HPT):

Acute encephalitis  
Acute meningitis  
Acute poliomyelitis  
Acute infectious hepatitis  
Anthrax  
Botulism  
Brucellosis  
Cholera  
COVID-19  
Diphtheria  
Enteric fever (typhoid or paratyphoid fever)  
Food poisoning  
Haemolytic uraemic syndrome (HUS)  
Infectious bloody diarrhoea  
Invasive group A streptococcal disease and scarlet fever  
Legionnaires' disease  
Leprosy  
Malaria  
Measles  
Meningococcal septicaemia  
Mumps  
Plague  
Rabies  
Rubella  
SARS  
Smallpox  
Tetanus  
Tuberculosis  
Typhus  
Viral haemorrhagic fever (VHF)  
Whooping cough  
Yellow fever